



Optos*Advance™*Merge Studies



Contents

Purpose	1
Merge Studies	1

Purpose

The document will provide guidance on how to combine two studies into a single study. You may need to perform this action if you imaged the same patient twice in one day.

Users must have the Edit Patient Study privilege in Optos Advance to merge studies.

Merge Studies

- 1 | Login to Optos Advance.
- 2 | Select the **checkbox** beside the two studies you want to merge.
- 3 | **Right-click** one of the studies, then select **Merge Studies.**

	Scleral-Buckle, I	Load Studies	
	RPE-Degenerati	Send to Device	
	Retinal-Hole-Op	Copy Link	
	Chorioretinitis, I	Lock/Unlock	
	Shadow-of-Mitte	Merge Studies	
	Retinitis-Pigmer	Merge Patients	
	Retinal-Degener	Delete	
	Hemorrhages,	Delete Pre-processed Results	
	Cystoid-Macular	Download Study	
•	Choroidal-Nevu	Apply Policy File to Folder	

4 | Information for the selected studies is shown.

G-109979/3





Optos*Advance™*Merge Studies



Attribute	Value	Value		
Last Name	Scleral-Buckle	RPE-Degeneration		
First Name	Daytona-1	Daytona-1		
Middle Name				
Patient Id	1037-D1	1035-D1		
Issuer Of Patient Id				
Gender	F 🔻	F 🔻		
DOB	19700101	19700101		
Study Id				
Performed On	20151108	20151108		
Accession Number				
Referring Physician	Optos^Library^^^	Optos^Library^^^		
Description				
Merge to left Merge to right Cancel				

- 5 | To merge the studies into the study shown on the left, select Merge to left.
- 6 | To merge the studies into the study shown on the right, select **Merge to right.**
- 7 | Select **Continue** to finalize combining studies into one single study.



Optos UK/Europe +44 (0)1383 843350 ics@optos.com **Optos North America** 800 854 3039 usinfo@optos.com **Optos DACH** DE: 0800 72 36 805 AT: 0800 24 48 86 CH: 0800 55 87 39 ics@optos.com **Optos Australia** +618 8444 6500 auinfo@optos.com 

